

## Package leaflet: Information for the patient

### [PRODUCT NAME] 250 mg tablets [PRODUCT NAME] 500 mg film-coated tablets abiraterone acetate

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What [PRODUCT NAME] is and what it is used for
2. What you need to know before you take [PRODUCT NAME]
3. How to take [PRODUCT NAME]
4. Possible side effects
5. How to store [PRODUCT NAME]
6. Contents of the pack and other information

#### **1. What [PRODUCT NAME] is and what it is used for**

[PRODUCT NAME] contains a medicine called abiraterone acetate. It is used to treat prostate cancer in adult men that has spread to other parts of the body. [PRODUCT NAME] stops your body from making testosterone; this can slow the growth of prostate cancer.

When [PRODUCT NAME] is prescribed for the early stage of disease where it is still responding to hormone therapy, it is used with a treatment that lowers testosterone (androgen deprivation therapy).

When you take this medicine, your doctor will also prescribe another medicine called prednisone or prednisolone. This is to lower your chances of getting high blood pressure, having too much water in your body (fluid retention), or having reduced levels of a chemical known as potassium in your blood.

#### **2. What you need to know before you take [PRODUCT NAME]**

##### **Do not take [PRODUCT NAME]**

- if you are allergic to abiraterone acetate or any of the other ingredients of this medicine (listed in section 6).
- if you are a woman, especially if pregnant. [PRODUCT NAME] is for use in male patients only.
- if you have severe liver damage.
- in combination with Ra-223 (which is used to treat prostate cancer).

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking this medicine.

#### **Warnings and precautions**

Talk to your doctor or pharmacist before taking this medicine:

---

pal (common) Abiraterone medac 250 mg tablets/ Abiraterone medac 500 mg film-coated tablets

Date of common version: 01/2021

<not for print>

- if you have liver problems
- if you have been told you have high blood pressure or heart failure or low blood potassium (low blood potassium may increase the risk of heart rhythm problems)
- if you have had other heart or blood vessel problems
- if you have an irregular or rapid heart rate
- if you have shortness of breath
- if you have gained weight rapidly
- if you have swelling in the feet, ankles, or legs
- if you have taken a medicine known as ketoconazole in the past for prostate cancer
- about the need to take this medicine with prednisone or prednisolone
- about possible effects on your bones
- if you have high blood sugar.

Tell your doctor if you have been told you have any heart or blood vessel conditions, including heart rhythm problems (arrhythmia), or are being treated with medicines for these conditions.

Tell your doctor if you have yellowing of the skin or eyes, darkening of the urine, or severe nausea or vomiting, as these could be signs or symptoms of liver problems. Rarely, failure of the liver to function (called acute liver failure) may occur, which can lead to death.

Decrease in red blood cells, reduced sex drive (libido), muscle weakness and/or muscle pain may occur.

[PRODUCT NAME] must not be given in combination with Ra-223 due to a possible increase in the risk of bone fracture or death.

If you plan to take Ra-223 following treatment with [PRODUCT NAME] and prednisone/prednisolone, you must wait 5 days before starting treatment with Ra-223.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking this medicine.

### **Blood monitoring**

[PRODUCT NAME] may affect your liver and you may not have any symptoms. When you are taking this medicine, your doctor will check your blood periodically to look for any effects on your liver.

### **Children and adolescents**

This medicine is not for use in children and adolescents. If [PRODUCT NAME] is accidentally ingested by a child or adolescent, go to the hospital immediately and take the package leaflet with you to show to the emergency doctor.

### **Other medicines and [PRODUCT NAME]**

Ask your doctor or pharmacist for advice before taking any medicine.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is important because [PRODUCT NAME] may increase the effects of a number of medicines including heart medicines, tranquilisers, some medicines for diabetes, herbal medicines (e.g., St John's wort) and others. Your doctor may want to change the dose of these medicines. Also, some medicines may increase or decrease the effects of [PRODUCT NAME]. This may lead to side effects or to [PRODUCT NAME] not working as well as it should.

Androgen deprivation treatment may increase the risk of heart rhythm problems. Tell your doctor if you are receiving medicine

- used to treat heart rhythm problems (e.g. quinidine, procainamide, amiodarone and sotalol);

---

pal (common) Abiraterone medac 250 mg tablets/ Abiraterone medac 500 mg film-coated tablets

Date of common version: 01/2021

<not for print>

- known to increase the risk of heart rhythm problems [e.g. methadone (used for pain relief and part of drug addiction detoxification), moxifloxacin (an antibiotic), antipsychotics (used for serious mental illnesses)].

Tell your doctor if you are taking any of the medicines listed above.

#### **[PRODUCT NAME] with food**

This medicine must not be taken with food (see section 3, “Taking this medicine”).

Taking [PRODUCT NAME] with food may cause side effects.

#### **Pregnancy and breast-feeding**

[PRODUCT NAME] is not for use in women. This medicine may cause harm to the unborn child if it is taken by women who are pregnant.

If you are having sex with a woman who can become pregnant, use a condom and another effective birth control method.

If you are having sex with a pregnant woman, use a condom to protect the unborn child.

<only for 250 mg>

[PRODUCT NAME] 250 mg tablets:

Women who are pregnant or who may be pregnant should wear gloves if they need to touch or handle [PRODUCT NAME].

#### **Driving and using machines**

This medicine is not likely to affect your being able to drive and use any tools or machines.

#### **[PRODUCT NAME] contains lactose and sodium**

- [PRODUCT NAME] contains lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicinal product contains 23.2 mg sodium (main component of cooking/table salt) per dose of four tablets for 250 mg and 23 mg sodium per dose of two tablets for 500 mg. This is equivalent to 1.2% of the recommended maximum daily dietary intake of sodium for an adult.

### **3. How to take [PRODUCT NAME]**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### **How much to take**

The recommended dose is 1,000 mg (four tablets) once a day.

The recommended dose is 1,000 mg (two tablets) once a day.

#### **Taking this medicine**

- Take this medicine by mouth.
- **Do not take [PRODUCT NAME] with food.**  
**Take [PRODUCT NAME] at least one hour before or at least two hours after eating** (see section 2, “[PRODUCT NAME] with food”).
- Swallow the tablets whole with water.
- Do not break the tablets.
- [PRODUCT NAME] is taken with a medicine called prednisone or prednisolone. Take the prednisone or prednisolone exactly as your doctor has told you.
- You need to take prednisone or prednisolone every day while you are taking [PRODUCT NAME].
- The amount of prednisone or prednisolone you take may need to change if you have a

---

pal (common) Abiraterone medac 250 mg tablets/ Abiraterone medac 500 mg film-coated tablets

Date of common version: 01/2021

<not for print>

medical emergency. Your doctor will tell you if you need to change the amount of prednisone or prednisolone you take. Do not stop taking prednisone or prednisolone unless your doctor tells you to.

Your doctor may also prescribe other medicines while you are taking [PRODUCT NAME] and prednisone or prednisolone.

**If you take more [PRODUCT NAME] than you should**

If you take more than you should, talk to your doctor or go to a hospital immediately.

**If you forget to take [PRODUCT NAME]**

If you forget to take [PRODUCT NAME] or prednisone or prednisolone, take your usual dose the following day.

If you forget to take [PRODUCT NAME] or prednisone or prednisolone for more than one day, talk to your doctor without delay.

**If you stop taking [PRODUCT NAME]**

Do not stop taking [PRODUCT NAME] or prednisone or prednisolone unless your doctor tells you to. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop taking [PRODUCT NAME] and see a doctor immediately if you notice any of the following:**

- Muscle weakness, muscle twitches or a pounding heart beat (palpitations). These may be signs that the level of potassium in your blood is low.

**Other side effects include:**

**Very common** (may affect more than 1 in 10 people):

Fluid in your legs or feet, low blood potassium, liver function test increases, high blood pressure, urinary tract infection, diarrhoea.

**Common** (may affect up to 1 in 10 people):

High fat levels in your blood, chest pain, irregular heart beat (atrial fibrillation), heart failure, rapid heart rate, severe infections called sepsis, bone fractures, indigestion, blood in urine, rash.

**Uncommon** (may affect up to 1 in 100 people):

Adrenal gland problems (related to salt and water problems), abnormal heart rhythm (arrhythmia), muscle weakness and/or muscle pain.

**Rare** (may affect up to 1 in 1,000 people):

Lung irritation (also called allergic alveolitis). Failure of the liver to function (also called acute liver failure).

**Not known** (frequency cannot be estimated from the available data): Heart attack, changes in ECG - electrocardiogram (QT prolongation), and serious allergic reactions with difficulty swallowing or breathing, swollen face, lips, tongue or throat, or an itchy rash.

Bone loss may occur in men treated for prostate cancer. [PRODUCT NAME] in combination with prednisone or prednisolone may increase bone loss.

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on

the safety of this medicine.

## 5. How to store [PRODUCT NAME]

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and the bottle label after EXP. The expiry date refers to the last day of that month.
- Do not use this medicine after the expiry date which is stated on the carton and the blister after EXP. The expiry date refers to the last day of that month.
- For 250 mg tablets: This medicine does not require any special temperature storage conditions. Store in the original package in order to protect from light.
- For 500 mg tablets: This medicine does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What [PRODUCT NAME] contains

- The active substance is abiraterone acetate. Each tablet contains 250 mg abiraterone acetate.
- The other ingredients are microcrystalline cellulose, croscarmellose sodium, lactose monohydrate, magnesium stearate, hypromellose (type 2910), Silica, colloidal anhydrous and sodium laurilsulfate (see section 2, “[PRODUCT NAME] contains lactose and sodium”).
- The active substance is abiraterone acetate. Each film-coated tablet contains 500 mg abiraterone acetate.
- The other ingredients are microcrystalline cellulose, croscarmellose sodium, lactose monohydrate, magnesium stearate, hypromellose (type 2910), Silica, colloidal anhydrous, sodium laurilsulfate (see section 2, “[PRODUCT NAME] contains lactose and sodium”). The film-coating contains iron oxide black (E172), iron oxide red (E172), Macrogol poly(vinyl alcohol) grafted copolymer, talc and titanium dioxide (E171).

### What [PRODUCT NAME] looks like and contents of the pack

- [PRODUCT NAME] tablets are white to off-white, oval, biconvex, uncoated tablets debossed with “250” on one side and plain on the other side.
- The tablets are provided in a HDPE bottle 100 ml with a child-resistant and desiccant including cap (polypropylene, PP). Each bottle contains 120 tablets. Each carton contains one bottle.
- [PRODUCT NAME] film-coated tablets are violet, oval, biconvex, film-coated tablets, debossed with “500” on one side and plain on the other side.
- The tablets are packaged in blisters in carton boxes containing 56 or 60 film-coated tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

[To be completed nationally]

### Manufacturer

[To be completed nationally]

**This medicinal product is authorised in the Member States of the EEA under the following names:**

---

pal (common) Abiraterone medac 250 mg tablets/ Abiraterone medac 500 mg film-coated tablets

Date of common version: 01/2021

<not for print>

[To be completed nationally]

**This leaflet was last revised in**